Section:	Division of Nursing	*******	Index:	6160.008a
	3	PROCEDURE	Page:	1 of 1
Approval:		***********	Issue Date: Revised Date:	August 14, 1997 January 2010
	HACKETTSTOWN REGIONAL MEDICAL CENTER			

Originator: P. Carpenito, RNC Revised by: P. Swanson, RN, MSN

**MATERNAL SERVICES** 

(Scope)

TITLE: EMLA CREAM

PURPOSE: To outline the procedure to use Emla cream to relieve discomfort at the site of insertion of IV or

epidural catheter.

SUPPORTIVE DATA: Emla cream can be used for all IV starts, for all blood work and for all insertions of epidural anesthesia

on all OB-GYN patients requiring a non-emergent IV, non-emergent lab work and/or non-emergent

epidural anesthesia.

Use requires a provider's order.

## CONTENT

## **PROCEDURE**

## **KEY POINTS**

**TECNNIQUE** 

 At Pyxis machine, enter patient's name and withdraw one five-gram tube of emla cream and obtain occlusive dressings for five areas. Fax order to pharmacy for profiling in pyxis.

Emla cream is indicated as a topical anesthetic for use on normal intact skin for local analgesia.

- 2. Determine two probable IV sites.
- 3. Prepare both antecubital areas (for lab work access)
- For epidural,(obtain order from anesthesia) determine space between L2 and L5 vertebrae (with patient lying on her side, palpate the anterior-superior iliac spine process, slide your hand to her back, this is approximately = to L4).

Cover the emla cream so that you get a thick layer underneath. Do not spread out the cream. Mark time of application directly on the occlusive dressing.

- Scan patient and medication using C5 for Cerner electronic MAR.
- 6. Use proper hand washing and use of gloves.
- Place a "dime" size amount of emla cream on selected sites and cover with an occlusive dressing. This must be done ½ - 2 hours before procedure is to be done.
- When ready to access site, simply remove dressing and proceed as usual.
- 9. Document in electronic nurse's notes.
- 10. Observe skin for any signs of local irritation.

**CONTRAINDICATIONS:** Emla cream (lidocaine 2.5% and prilocaine 2.5%) is contraindicated in patients with a known history of sensitivity to local anesthetics of the amide type or to any other component of the product.

Reference: Product insert: Astra\* Westborough, Ma 01581, Lycka, Ba Medline Record